

# Ising Expansion for the Hubbard Model

Zhu-Pei Shi and Rajiv R. P. Singh

*Department of Physics, University of California, Davis, California 9561 6*

## Abstract

We develop series expansions for the ground state properties of the Hubbard model, by introducing an Ising anisotropy into the Hamiltonian. For the two-dimensional (2D) square lattice half-filled Hubbard model, the ground state energy, local moment, sublattice magnetization, uniform magnetic susceptibility and spin stiffness are calculated as a function of  $U/t$ , where  $U$  is the Coulomb constant and  $t$  is the hopping parameter. Magnetic susceptibility data indicate a crossover around  $U \approx 4$  between spin density wave antiferromagnetism and Heisenberg antiferromagnetism. Comparisons with Monte Carlo simulations, RPA result and mean field solutions are also made.

PACS numbers: 71.27.+a, 75.10.Jm, 75.40.Cs

## I. INTRODUCTION

Low temperature properties of strongly correlated electron systems have been studied extensively over the past few decades. The Hubbard model is one of the simplest models of interacting electrons, which is believed to capture the main features of electron correlations in condensed matter systems. The model is best understood at half filling, where it exhibits the Mott-Hubbard phenomena, namely the existence of insulating behavior accompanied by magnetic order in a half-filled band of electrons. Recent discovery of high- $T_c$  superconductivity in doped antiferromagnets (copper oxides) has led Anderson [1] to suggest that superconductivity in these materials is closely related to the physics of strong correlations contained in the Hubbard model. The spin-half Heisenberg model (a large  $U$  limit of the Hubbard model) has formed the basis for a quantitative understanding of the magnetic excitations in the insulating stoichiometric phase of the high- $T_c$  materials. Studies of the Hubbard and  $t - J$  models have also provided qualitative insights into the magnetic and transport properties of doped high- $T_c$  materials [2].

The Hubbard model is defined by the lattice Hamiltonian

$$H = -t \sum_{\langle i,j \rangle, \sigma} (c_{i\sigma}^+ c_{j\sigma} + c_{j\sigma}^+ c_{i\sigma}) + U \sum_i (n_{i\uparrow} - \frac{1}{2})(n_{i\downarrow} - \frac{1}{2}) - \mu \sum_i (n_{i\uparrow} + n_{i\downarrow}) \quad , \quad (1)$$

where, as usual,  $c_{i\sigma}^+$  and  $c_{i\sigma}$  are the creation and annihilation operators for electrons with a  $z$ -component of spin  $\sigma$  at lattice site  $i$ , and  $n_{i,\sigma} = c_{i\sigma}^+ c_{i\sigma}$ .  $U$  is the on site repulsive interaction,  $\mu$  the chemical potential, and  $t$  the nearest-neighbor hopping amplitude. The sum  $\langle i, j \rangle$  is over all pairs of nearest neighbor lattice sites.

In spite of its apparent simplicity, the Hubbard model is exactly soluble only in one-dimension [3]. In two-dimensions, relevant to the high- $T_c$  materials, this model has been studied by a variety of approximate methods. These include finite size diagonalization [2], mean-field theory [4], Green's-function decoupling schemes [5], variational approaches [6], the random phase approximation (RPA) [7], high temperature series expansion [8] and quantum Monte Carlo simulations [9–11]. Despite these efforts some of the very basic features of this

model, such as its phase diagram in the space of temperature, density and  $U/t$  are not fully known. The numerical studies do provide a consistent picture for the magnetically ordered zero temperature phase at half-filling. However, away from half-filling the possibility of long range incommensurate magnetic order and superconductivity remain open.

In this paper we develop a series expansion method for studying the Hubbard model, and calculate the ground state energy  $E_g$ , local moment  $L$ , sublattice magnetization  $M^+$ , the uniform magnetic susceptibility  $\chi_\perp$  and spin stiffness  $\rho_s$ , for the half-filled case, as a function of  $U/t$ . This study is complimentary to the existing numerical methods and provides additional insight into this model. Our results are in broad agreement with the Monte Carlo simulations. There are however, some quantitative differences. For example the manner in which the local moment  $L$  goes to  $1/2$  as  $U \rightarrow 0$ , is different in our calculations than in the Monte-Carlo simulations. Our results are much closer to the spin-density-wave mean field results in this limit. Furthermore, the behavior of magnetic susceptibility as a function of  $U/t$  changes character at  $U/t \approx 4$  indicating a crossover between spin density wave (SDW) antiferromagnet and a Heisenberg antiferromagnet at this value

The organization of the paper is as follows: In Sec. II we discuss our series expansion method for the two-dimensional Hubbard model. In Sec. III we present results for the ground state properties and in Sec. IV for the uniform magnetic susceptibility of the model. In Sec. V we derive and compute the spin stiffness of the Hubbard model and estimate the spin wave velocity from the relation  $v_s^2 = \rho_s/\chi_\perp$ . Finally, in Sec. VI, we summarize our results.

## II. ISING EXPANSION FOR THE HUBBARD MODEL

In order to develop series expansion for the Hubbard model, we need to introduce an Ising anisotropy into the Hubbard Hamiltonian:

$$H_\lambda = -\lambda t \sum_{\langle i,j \rangle, \sigma} (c_{i\sigma}^+ c_{j\sigma} + c_{j\sigma}^+ c_{i\sigma}) + U \sum_i (n_{i\uparrow} - \frac{1}{2})(n_{i\downarrow} - \frac{1}{2}) + f(\lambda) \sum_{\langle i,j \rangle} \sigma_i^z \sigma_j^z \quad , \quad (2)$$

where  $\sigma_i^z = (n_{i\uparrow} - n_{i\downarrow})$  is the  $z$  component of the spin at site  $i$ . The particle-hole symmetry ensures half-filling. The function  $f(\lambda)$  needs to be chosen such that  $f(0 \leq \lambda < 1) > 0$ , and  $f(1) = 0$ . For  $\lambda = 0$  the atomic limit of the Hubbard model is highly degenerate, however the Ising term selects from these the Néel states as the two degenerate ground states. Furthermore, this term also introduces a gap in the spectrum at  $\lambda = 0$ . *For  $\lambda = 1$  the Ising anisotropy goes to zero and the conventional Hubbard Hamiltonian is recovered.* Ground state properties of the model for  $\lambda \neq 1$  can be obtained by an expansion in powers of  $\lambda$ . If the gap does not close before  $\lambda = 1$  as expected for this model, we can obtain properties of the Hubbard model by extrapolating the expansions to  $\lambda = 1$ .

We choose the function  $f(\lambda)$  to have the form

$$f(\lambda) = J(1 - \lambda) \quad , \quad (3)$$

where  $J$  is a constant which can be tuned to improve the convergence of the extrapolations. In the strong coupling limit, the half-filled Hubbard model is equivalent to an antiferromagnetic Heisenberg model with Hamiltonian:

$$H_{Heisenberg} = J_H \sum_{\langle i,j \rangle} (\sigma_i \cdot \sigma_j - 1) \quad , \quad (4)$$

where  $\langle i, j \rangle$  denotes nearest neighbors and  $J_H = t^2/U$ . If we choose  $J$  in Eq.(3) to be much large than  $J_H$  we find that the expansion coefficients become very large and oscillate rapidly, making extrapolations very difficult. On the other hand, if  $J$  is much smaller than  $J_H$ , the expansion coefficients become very small and a large number of terms will be needed to extrapolate to  $\lambda = 1$ . The optimum value of  $J$  is around  $J_H$ .

In order to employ series expansion techniques, we write the Hamiltonian in Eq.(2) as

$$H_\lambda = H_0 + H' \quad , \quad (5)$$

where unperturbed part of Hamiltonian  $H_0$  is defined as

$$H_0 = U \sum_i (n_{i\uparrow} - \frac{1}{2})(n_{i\downarrow} - \frac{1}{2}) + J \sum_{\langle i,j \rangle} \sigma_i^z \sigma_j^z \quad , \quad (6)$$

and the perturbation  $H'$  is

$$H' = -\lambda t \sum_{\langle i,j \rangle, \sigma} (c_{i\sigma}^+ c_{j\sigma} + c_{j\sigma}^+ c_{i\sigma}) - \lambda J \sum_{\langle i,j \rangle} \sigma_i^z \sigma_j^z . \quad (7)$$

Throughout this paper, we take  $t = 1$  as the energy unit.  $U$  and  $J$  are two interaction parameters. The parameter  $\lambda$  simply allows us to extrapolate from the Ising model to the half-filled Hubbard model.

### III. GROUND STATE PROPERTIES

The ground state energy  $E_g$  is defined as

$$\begin{aligned} E_g &= \langle \psi_g(\lambda) | H(\lambda) | \psi_g(\lambda) \rangle / \langle \psi_g(\lambda) | \psi_g(\lambda) \rangle \\ &= \sum_n e_n(U, J) \lambda^n , \end{aligned} \quad (8)$$

where  $|\psi_g(\lambda)\rangle$  is the ground state.

The sublattice magnetization  $M^+$  is defined as

$$\begin{aligned} 2M^+ &= \frac{1}{N} \langle \sum_i \epsilon(i) (n_{i\uparrow} - n_{i\downarrow}) \rangle \\ &= \sum_n m_n(U, J) \lambda^n , \end{aligned} \quad (9)$$

where  $N$  is the number of lattice sites,  $\epsilon(i)$  set to be  $+1$  on one sublattice and  $-1$  on the other, and the angular brackets refer to ground state expectation values.

The squared local moment  $L$  is defined via the relation

$$\begin{aligned} L &= \frac{1}{N} \langle \sum_i (n_{i\uparrow} - n_{i\downarrow})^2 \rangle \\ &= \sum_n l_n(U, J) \lambda^n . \end{aligned} \quad (10)$$

The expansion coefficients for  $E_g$ ,  $L$ , and  $2M^+$  to 11th order in  $\lambda$  are listed in tables I-III.

We find that properties which are not sensitive to long range order, such as ground state energy  $E_g$  and local moment  $L$  can be easily obtained to high accuracy with the simplest extrapolations. Padé approximants [12] to the function under consideration may be formed

directly. In order to estimate the values of  $E_g$  and  $L$  at particular  $U$ , we compute all Padé approximants from series up to 9th, 10th and 11th orders for three different Ising parameter  $J/(t^2/U) = 0.75, 1.0, 1.25$ , and remove ill-behaved Padé approximants. The estimates are set to be the median of the Padé approximants at  $\lambda = 1$ , and the errorbars are estimated as half the spread of these Padé approximants.

The values of the ground state energies at  $\lambda = 1$  are  $E_g = -1.51(2), -1.86(1), -2.934(4)$  and  $-5.2293(8)$  for  $U = 1, 4, 10$  and  $20$ , respectively. They are plotted in Fig. 1 as solid circles with errorbars. The solid line is a guide to the eye. Our calculated ground state energies  $E_g$  agree with the Monte Carlo simulations. For example, at  $U = 4$ , the extrapolated ground state energy from finite temperature Monte Carlo simulations [10] is  $-1.864(1)$ . Projected Monte Carlo simulations [13] (an algorithm which allows the calculation of ground state properties in the canonical ensemble, *i.e.*, with a fixed number of electrons) gives  $-1.864(2)$ . At  $U = 1$ , our result ( $E_g + U/4 = -1.26$ ) is close to the slave-boson mean field result ( $\approx -1.3$ ) [14]

The squared local moment obtained from the Padé approximants at  $\lambda = 1$  are  $L = 0.54(1), 0.75(1), 0.924(2)$  and  $0.9788(4)$  for  $U = 1, 4, 10$  and  $20$ , respectively. They are plotted in Fig. 2 as solid circles with errorbars. Once again, the solid line is a guide to the eye. The empty squares and dashed lines are the Monte Carlo simulations obtained by White *et al.* [10] at low temperatures [Fig. 6(a) in Ref. 9]. We see that both calculations are in good agreement with each other at large  $U$  but they show different behaviors at small  $U$ . Monte Carlo simulations show that local moment  $L$  increases rapidly as  $U$  increases (*concave function*) while ours indicate a relatively slow growth (*convex function*) at small  $U$  region, as shown in Fig. 3. We would naively expect the series extrapolation to be better at large  $U$ , and not so well at small  $U$ . However, as we now show, our results are rather close to the weak coupling results at small  $U$ , which suggest that our convergence is reasonable for all  $U$ .

Within the Hartree-Fock approximation, we can derive a simple relationship between local moment and sublattice magnetization. The local moment is

$$L = \langle n_{\uparrow} \rangle + \langle n_{\downarrow} \rangle - 2 \langle n_{\uparrow} \rangle \langle n_{\downarrow} \rangle , \quad (11)$$

where

$$\begin{aligned} \langle n_{\uparrow} \rangle &= 0.5 + M^+ \\ \langle n_{\downarrow} \rangle &= 0.5 - M^+ . \end{aligned} \quad (12)$$

Substituting Eq.(12) into Eq.(11) we have

$$L = 2 (M^+)^2 + 0.5 , \quad (13)$$

The mean field solution for the sublattice magnetization is [7,15]

$$M^+ = \Delta/U , \quad (14)$$

where  $\Delta$  can be obtained by solving the gap equation

$$\frac{U}{2N} \sum_{\mathbf{k}} \frac{1}{(\epsilon_{\mathbf{k}}^2 + \Delta^2)^{1/2}} = 1 . \quad (15)$$

$N$  is the total number of sites and  $\epsilon(\mathbf{k}) = -2t(\cos k_x + \cos k_y)$ . The local moment given by numerical solution of the gap equation is plotted in Fig. 2, a dashed line. One can see that the function  $L(U)$  has a convex shape at small  $U$  which agrees with our conclusion. At  $U = 1$ , our result ( $L = 0.54$ ) agrees with the slave-boson mean field result ( $L \approx 0.53$ ) [14]

The extrapolation of sublattice magnetization  $M^+$  appears much harder. In the 2D spin- $\frac{1}{2}$  square lattice Heisenberg antiferromagnet, the sublattice magnetization ( $2M^+$ ) is estimated by expansions around the Ising limit. In order to extrapolate the  $2M^+$  series reliably, one has to remove the singularity of the form  $[1 - (J_{\perp}/J_{\parallel})^2]^{\frac{1}{2}}$  caused by Goldstone modes [16], where  $J_{\perp}$  is the exchange perpendicular to the direction of ordering and  $J_{\parallel}$  the exchange along it. One way of doing this is to go a new variable  $\delta$  given by [17]

$$1 - \delta = [1 - (J_{\perp}/J_{\parallel})^2]^{\frac{1}{2}} . \quad (16)$$

We expect to have the same situation for the sublattice magnetization  $2M^+$  in the 2D Hubbard model because it is equivalent to the Heisenberg model at large  $U$ . Here we set a new variable  $\delta$  to be

$$1 - \delta = (1 - \lambda)^{\frac{1}{2}} \quad , \quad (17)$$

where  $\delta = 1$  at  $\lambda = 1$ . Then the series for the sublattice magnetization  $2M^+$  becomes

$$2M^+ = \sum_{n=0}^{11} b_n(U, J) \delta^n \quad , \quad (18)$$

where  $b_n(U, J)$  can be calculated from previous series  $m_n(U, J)$ . Again, we compute all Padé approximants from 7th order to 11th order series. The median of Padé approximants at  $\delta = 1$  give  $2M^+ = 0.12(5), 0.37(5), 0.58(3)$ , and  $0.61(3)$  for  $U = 1, 4, 10$  and  $20$ , respectively. They are plotted in Fig. 3, solid circles with errorbars. The solid line is a guide to the eye.

At large  $U$ , the effective Hamiltonian for the half-filled Hubbard model is known to be the spin- $\frac{1}{2}$  Heisenberg model. The quantum zero-point spin wave fluctuations reduce the sublattice magnetization of  $2M^+$  from the classical value of unity to about  $2M^+ = 0.6$  as first obtained by Anderson [18]. This is also supported by Monte Carlo simulations [19] and series expansion studies [16]. At  $U = 20$ , we estimate  $2M^+ = 0.61$  which is close to the value 0.6 of the Heisenberg model. Also, at  $U = 4$ , we can compare our results [ $2M^+ = 0.37(5)$ ] with the Monte Carlo simulations [ $2M^+ = 0.39(5)$ ] of Hirsch and Tang [9] and find fairly good agreement.

Also, we compare our results with the mean field solutions and the RPA calculations. The sublattice magnetization given by numerical solution of the gap equation is plotted in Fig. 3, as a long dashed line. At large  $U$ , the mean field solution gives  $2M^+ = 1$ , which is far from the true value ( $2M^+ = 0.6$ ) of the Heisenberg model because of the neglect of the quantum fluctuations. In order to take into account the fluctuation effects on the sublattice magnetization, one simply has to calculate the self-energy correction to the one particle Green's function as done by Schrieffer, Wen and Zhang [7] in random phase approximation. Their RPA result is plotted in Fig. 3 by a dotted line. Clearly, the RPA result brings the mean field result much closer to our numerical estimates.

We note again that the purpose of introducing the Ising anisotropy  $J(1 - \lambda)\sigma_i^z \sigma_j^z$  into the conventional Hubbard Hamiltonian is to be able to carry out the series expansions. The

ground state properties of the Hubbard model should not depend on the parameter  $J$  of the Ising anisotropy. One can also choose another type of function  $f(\lambda)$  in Eq.(2):

$$f(\lambda) = J (1 - \lambda^2) \quad . \quad (19)$$

This form of  $f(\lambda)$  leads to an expansion in powers of  $\lambda^2$ . This allows one to treat the  $J$  term generated by the perturbation in the Hubbard model and the Ising term introduced by hand on *an equal footing*. However, it has the disadvantage that the number of series coefficients available for extrapolation is reduced by half. The resulting extrapolations at  $\lambda = 1$  are consistent with those obtained earlier.

#### IV. MAGNETIC SUSCEPTIBILITY

We now discuss the behavior of the uniform magnetic susceptibility of the 2D Hubbard model from the series expansions. We turn on the magnetic field and add a Zeeman interaction into the 2D Hubbard Hamiltonian:

$$\begin{aligned} H_{\lambda h} = & -\lambda t \sum_{\langle i,j \rangle, \sigma} (c_{i\sigma}^+ c_{j\sigma} + c_{j\sigma}^+ c_{i\sigma}) + U \sum_i (n_{i\uparrow} - \frac{1}{2})(n_{i\downarrow} - \frac{1}{2}) \\ & + J(1 - \lambda) \sum_{\langle i,j \rangle} \sigma_i^z \sigma_j^z - h \sum_i \sigma_i^x \quad , \end{aligned} \quad (20)$$

where  $h$  is magnetic field and  $\frac{1}{2}\sigma_i^x$  is the  $x$ -component of the spin at site  $i$ . The uniform magnetic susceptibility  $\chi_{\perp}$  is defined as

$$\begin{aligned} 4\chi_{\perp} &= -\partial^2 E / \partial h^2 \big|_{h=0} \\ &= \sum_n x_n(U, J) \lambda^n \quad . \end{aligned} \quad (21)$$

The expansion coefficients of  $4\chi_{\perp}$  to 9th order in  $\lambda$  are given in Table IV.

At half-filling, 2D Hubbard model has antiferromagnetic long range order. One can see that  $\chi_{\perp}$  series is dominated by a simple pole at  $\lambda = -1$ . We need to remove this singularity before further analysis. This is done by going to a new variable  $\beta$  given by

$$\beta = 2\lambda / (1 + \lambda) \quad . \quad (22)$$

The resulting series has the form

$$4\chi_{\perp} = \sum_{n=0}^9 y_n(U, J) \beta^n, \quad (23)$$

where  $y_n(U, J)$  can be obtained from the previous series  $x_n(U, J)$ . But this new series still contains singularities at the Hubbard point ( $\lambda = 1$  or  $\beta = 1$ ). Going to a new variable  $\delta$  defined by

$$1 - \delta = (1 - \beta)^{1/2}, \quad (24)$$

and constructing Padé approximants, we find that only two Padé approximants ( $K = 4, M = 5$  and  $K = 5, M = 4$ ) from 9th order series show good behavior and both give the same results. For this reason it is difficult to estimate errors in these results. The uniform magnetic susceptibility obtained from these Padé approximants are  $4\chi_{\perp} = 0.94, 0.58, 0.75, 0.92$  and  $1.40$  for  $U = 1, 4, 6, 10$  and  $20$ , respectively. They are plotted in Fig. 4 as solid circles. The solid line is a guide to the eye.

At  $U = 20$ ,  $\chi_{\perp}/U = \chi_{\perp} J = 0.0175$  is close to the large  $U$  limit value  $\chi_{\perp} J_H = 0.0163$  of the Heisenberg model [16] shown by a dashed line in Fig. 4(a). At  $U = 4$ , our result ( $4\chi_{\perp} = 0.58$ ) agrees with the estimated value ( $\approx 0.53$ ) of Monte Carlo simulations at low temperature by White *et al.* [10], but does not agree with the estimated value ( $\approx 1.0$ ) of projected Monte Carlo simulations by Furukawa and Imada [13]. At  $U = 10$ , our result ( $4\chi_{\perp} = 0.92$ ) agrees with the estimated value ( $\approx 0.98$ ) of Monte Carlo simulations by Moreo [20].

One can see that our magnetic susceptibility data indicate a dip at  $U = U_c \approx 4$ , as shown in the Fig. 4(b). It suggests that there is a crossover around  $U_m$  in the behavior of the 2D Hubbard model at half filling. For  $U > U_c$ , we can see that magnetic susceptibility ( $4\chi_{\perp}$ ) is almost proportional to  $U$  or inversely proportional to the spin superexchange  $J$ . This can be taken as evidence that the magnetic state for  $U > U_c$  for 2D Hubbard model is that of the Heisenberg model. But, for  $U < U_c$ , magnetic susceptibility ( $4\chi_{\perp}$ ) has a different behavior, and decreases as  $U$  increases. This can be interpreted as a weak coupling behavior of the

SDW ground state. Thus we obtain a crossover between these types of behavior around  $U \approx 4$ .

Also, we compare our result with the mean field solutions of uniform magnetic susceptibility [21]:

$$\chi_{\perp} = \frac{1}{2U} [(1 - \Delta^2 U I_1 - U I_2^2/I_1)^{-1} ; -1] \quad , \quad (25)$$

where

$$I_1 = \frac{1}{2N} \sum_{\mathbf{k}} \frac{1}{E^3(\mathbf{k})}$$

$$I_2 = -\frac{1}{2N} \sum_{\mathbf{k}} \frac{\epsilon(\mathbf{k})}{E^3(\mathbf{k})} \quad .$$

Here  $E(\mathbf{k}) = [\epsilon^2(\mathbf{k}) + \Delta^2]^{1/2}$ . It is plotted in Fig. 4(b), a solid line. We see that the mean field result is qualitatively correct, but overestimates  $\chi_{\perp}$ , especially at large  $U$ .

## V. SPIN STIFFNESS OF THE HUBBARD MODEL

The spin stiffness constant  $\rho_s$  is a measure of the response of the spin system in an ordered phase to a twist of the order parameter. To our knowledge, there are no quantum Monte Carlo estimates for the spin stiffness in the Hubbard model. Derivation and calculation of this quantity for the Hubbard model was discussed by us recently in a short communication [22]. For completeness, we discuss it again here.

If we rotate the ordering direction by a small angle  $\theta$  along a given direction such as  $y$  axis, then the spin stiffness constant  $\rho_s$  can be defined through the increase of the ground state energy:

$$E_g(\theta) = E_g(\theta = 0) + \frac{1}{2} \rho_s \theta^2 + O(\theta^4) \quad , \quad (26)$$

This rotation can be carried out by the following transformation applied to the Fermion operators:

$$\begin{pmatrix} c'_{\uparrow} \\ c'_{\downarrow} \end{pmatrix} = \begin{pmatrix} \cos \phi/2 & \sin \phi/2 \\ -\sin \phi/2 & \cos \phi/2 \end{pmatrix} \begin{pmatrix} c_{\uparrow} \\ c_{\downarrow} \end{pmatrix} \quad , \quad (27)$$

and

$$\begin{pmatrix} c_{\uparrow}'^+ \\ c_{\downarrow}'^+ \end{pmatrix} = \begin{pmatrix} \cos \phi/2 & \sin \phi/2 \\ -\sin \phi/2 & \cos \phi/2 \end{pmatrix} \begin{pmatrix} c_{\uparrow}^+ \\ c_{\downarrow}^+ \end{pmatrix} . \quad (28)$$

This transformation generates desired spin rotation:

$$\begin{aligned} \sigma_{z'} &= c_{\uparrow}'^+ c_{\uparrow}' - c_{\downarrow}'^+ c_{\downarrow}' \\ &= \sigma_z \cos \phi + \sigma_x \sin \phi \\ \sigma_{x'} &= c_{\uparrow}'^+ c_{\downarrow}' - c_{\downarrow}'^+ c_{\uparrow}' \\ &= -\sigma_z \sin \phi + \sigma_x \cos \phi \end{aligned} \quad (29)$$

After rotation by a *relative* angle  $\theta$  ( that is letting  $\phi$  change by  $\theta/2$  ) between neighboring sites separated along  $y$  axis ( $\hat{y}$  denotes unit distance in  $y$  direction) ,  $H$  in Eq.(1) becomes

$$H_{\theta} = H + H^{dia} + H^{para} + O(\theta^3) , \quad (30)$$

where

$$\begin{aligned} H^{dia} &= \frac{t\theta^2}{8} \sum_{i,\sigma} (c_{i\sigma}^+ c_{i+\hat{y}\sigma} + c_{i+\hat{y}\sigma}^+ c_{i\sigma}) \\ H^{para} &= -\frac{t\theta}{2} \sum_i (c_{i\uparrow}^+ c_{i+\hat{y}\downarrow} - c_{i\downarrow}^+ c_{i+\hat{y}\uparrow} \\ &\quad + c_{i+\hat{y}\downarrow}^+ c_{i\uparrow} - c_{i+\hat{y}\uparrow}^+ c_{i\downarrow}) . \end{aligned} \quad (31)$$

The “diamagnetic” term  $H^{dia}$  is already of order  $\theta^2$  so for the calculation of the energy to order  $\theta^2$ , it can be replaced by its expectation value in the ground state of the  $\theta = 0$  Hamiltonian. We have

$$\begin{aligned} \rho_s^{dia} &= -\frac{1}{8} (Kinetic Energy) \\ &= -\frac{1}{8} [E_g(\theta = 0) - \frac{U}{2} (n - L)] , \end{aligned} \quad (32)$$

where  $n$  is a band filling, and  $L$  a local moment defined in Eq.(10). We have already obtained  $E_g$  and  $L$  in Sec. III, and  $n = 1$  for half-filling.

The contribution of the “paramagnetic” term  $H^{para}$  to the ground state energy in order  $\theta^2$  can be obtained by treating it in second order perturbation theory. We have

$$\begin{aligned}\rho_s^{para} &= 2 \frac{\partial^2 E}{\partial \theta^2} \Big|_{\theta=0} ; \\ &= \sum_n p_n \lambda^n ,\end{aligned}\tag{33}$$

where  $E$  represents energy of the Hamiltonian  $H_\lambda + H^{para}$ . The series coefficients  $p_n$  of  $\rho_s^{para}$  to 9th order in  $\lambda$  are given in the Table V.

We use the same Padé analysis for  $\rho_s^{para}$  as for uniform magnetic susceptibility. We obtain the spin stiffness  $\rho_s = \rho_s^{dia} + \rho_s^{para} = 0.186(15), 0.15(1), 0.077(7)$  and  $0.039(5)$  for  $U = 1, 4, 10$  and  $20$ , respectively. They are plotted in Fig. 5 as filled circles with a dashed line as guide to the eye. The errorbars represent the spread in the Padé estimates. At  $U = 20$ , our result ( $\rho_s U = 0.78$ ) agrees well with the value (0.73) of the Heisenberg model [16] shown by a short solid line.

We also compare our results with the Hartree-Fock approximation for the spin stiffness [21]:

$$\rho_s = -\frac{t}{2N} \sum_{\mathbf{k}} \frac{\epsilon(\mathbf{k}) \cos k_x}{E(\mathbf{k})} .\tag{34}$$

The spin stiffness given by mean field solution of the gap equation is plotted in Fig. 5(b) as a solid line. One can see that the mean field approximation overestimates the stiffness at large  $U$ . At  $U = 20$ , the mean field result is  $\rho_s U = 0.98$  compared to 0.73 for the Heisenberg model.

We note that the spin stiffness (filled squares in Fig. 1) from the variational Monte Carlo method with a Gutzwiller-type wave function are even larger than values of the mean field solution [21]. We believe that the large discrepancy is due to the missing spin flip processes in the Gutzwiller variational wave function used in [21]. Their calculations only get contributions to the spin-stiffness beyond the Hartree-Fock result from the “diamagnetic” term, whereas the “paramagnetic” part of the spin stiffness which contains spin-flip processes does not get corrected. The expectation value of the operator  $H^{para}$  between the Hartree-Fock and Gutzwiller wave functions is zero [23]. This suggests that one might improve the

variational Monte Carlo method by adding a spin flip operator  $O_{sp}$  into the Gutzwiller-type wave function:

$$| \Psi_{GW} \rangle = \prod_i [1 - (1 - g)n_{i\uparrow} n_{i\downarrow} + p \times O_{sp}] | \Psi_{HF} \rangle$$

where  $p$  is a parameter and

$$O_{sp} = (c_{i\uparrow}^\dagger c_{i+\hat{y}\downarrow} - c_{i\downarrow}^\dagger c_{i+\hat{y}\uparrow} + c_{i+\hat{y}\downarrow}^\dagger c_{i\uparrow} - c_{i+\hat{y}\uparrow}^\dagger c_{i\downarrow}) \quad .$$

We now turn to the calculation of spin-wave velocity. This can be calculated analytically by adding RPA fluctuations to the SDW state [7]. Designed, at first sight, for intermediate and weak coupling ( $U/4t \leq 1$ ) this method also interpolates smoothly to the Heisenberg limit at large  $U/t$ .

Alternatively, there is a relationship between the spin stiffness  $\rho_s$  of the order parameter with respect to a spiral twist, the spin wave velocity  $v_s$  and the uniform magnetic susceptibility  $\chi_\perp$  transverse to the staggered spin density:

$$\rho_s = v_s^2 \chi_\perp \quad . \quad (35)$$

It was originally derived by the hydrodynamic theory of Halperin and Hohenberg [24]. The spin wave velocity obtained from this relationship ( $\rho_s = v_s^2 \chi_\perp$ ) are  $v_s = 0.89, 1.02, 0.58$  and  $0.34$  for  $U = 1, 4, 10$  and  $20$ , respectively. They are plotted in Fig. 6 as filled circles with a solid line as a guide to the eye. At  $U = 20$ , our  $v_s = 0.34$  agrees well with the limiting value ( $v_s = 1.18\sqrt{2}J' = 0.33$ ) of the Heisenberg model [16], a dashed line in Fig. 6, where  $J' = 4t^2/U$ .

We compare this result with the RPA solution of Schrieffer, Wen and Zhang [7]:

$$v_s = [4t^2(1 - \Delta^2 U I_1) \frac{I_3}{I_1}]^{1/2} \quad , \quad (36)$$

where

$$I_3 = \frac{1}{2N} \sum_{\mathbf{k}} \frac{\sin^2(k_x)}{E^3(\mathbf{k})} \quad . \quad (37)$$

It is plotted in Fig. 6 as a dotted line. We see that the RPA result for the spin wave velocity are lower than our series expansion results.

## VI. SUMMARY

To summarize, in this paper we developed series expansions for the half-filled Hubbard model, by introducing an Ising anisotropy into the conventional Hubbard Hamiltonian. Series were developed for the ground state energy, the local moment, the sublattice magnetization, the uniform magnetic susceptibility and the spin stiffness, and were extrapolated to the Hubbard model using standard series extrapolation methods. In general, our results are in good agreement with the quantum Monte Carlo simulations. There are some quantitative differences, which are also discussed. Our magnetic susceptibility data indicate a crossover between SDW antiferromagnetic state at small  $U$  and a Heisenberg antiferromagnet at large  $U$  at  $U \approx 4$ . We note that our results for  $\rho_s$  as well as  $M^\dagger$  are somewhat higher than the known results for the Heisenberg models. We believe this reflects the fact that the series have not converged as well as for the Heisenberg model and hence the reduction in these quantities due to the zero-point spin-wave fluctuations is not fully accounted for. Still, the convergence ( $\sim 5\%$ ) is quite reasonable.

This research was supported in part by NSF (DMR-9318537) and University Research Funds of the University of California at Davis. Zhu-Pei Shi would like to thank Prof. Richard T. Scalettar and Prof. Barry M. Klein for discussions.

## REFERENCES

- [1] P. W. Anderson, *Frontiers and Borderlines in Many-Particle Physics*, edited by R. A. Broglia and J. R. Schrieffer (North-Holland, Amsterdam, 1988).
- [2] E. Dagotto, Rev. Mod. Phys. **66**, 763 (1994), and references therein.
- [3] E. Lieb and F. Wu, Phys. Rev. Lett. **20**, 1445 (1968).
- [4] D. Penn, Phys. Rev. **142**, 350 (1966).
- [5] J. Hubbard, Proc. R. Soc. London, Ser. A **276**, 283 (1963); **281**, 401 (1964).
- [6] M. C. Gutzwiller, Phys. Rev. **137**, A1726 (1965).
- [7] J. R. Schrieffer, X. G. Wen and S. C. Zhang, Phys. Rev. B **39**, 11663 (1989).
- [8] Y. S. Yang and C. J. Thompson, J. Phys. A **24**, L279 (1991); J. A. Henderson, J. Oitmaa and M. C. B. Asley, Phys. Rev. B **46**, 6328 (1992).
- [9] J. E. Hirsch and S. Tang, Phys. Rev. Lett. **62**, 591 (1989).
- [10] S. R. White, D. J. Scalapino, R. L. Sugar, E. Y. Loh, J. E. Gubernatis, R. T. Scalettar, Phys. Rev. B **40**, 506 (1989).
- [11] A. Moreo, *et al.*, Phys. Rev. B **41**, 2313 (1990).
- [12] D. L. Hunter and G. A. Baker, Jr., Phys. Rev. B **19**, 3808 (1979).
- [13] N. Furukawa and K. Imada, J. Phys. Soc. Jpn. **61**, 3331 (1992).
- [14] L. Lilly, A. Muramatsu, and W. Hanke, Phys. Rev. Lett. **65**, 1379 (1990).
- [15] J. E. Hirsch, Phys. Rev. B **31**, 4403 (1985).
- [16] R. R. P. Singh, Phys. Rev. B **39**, 9760 (1989); R. R. P. Singh and D. A. Huse, Phys. Rev. B **40**, 7247 (1989).
- [17] D. A. Huse, Phys. Rev. B **37**, 2380 (1988).

- [18] P. W. Anderson, Phys. Rev. **86**, 694 (1952).
- [19] J. D. Reger and A. P. Young, Phys. Rev. B **37**, 5978 (1988).
- [20] A. Moreo, Phys. Rev. B **48**, 3380 (1993).
- [21] P. J. H. Denteneer and J. M. J. van Leeuwen, Europhysics Lett. **20**, 413 (1993).
- [22] Zhu-Pei Shi and Rajiv R. P. Singh, to be published.
- [23] P. J. H. Denteneer, G. An and J. M. J. van Leeuwen, Phys. Rev. B **47**, 6256 (1993).
- [24] B. I. Halperin and P. C. Hohenberg, Phys. Rev. **188**, 898 (1969).

# TABLES

TABLE I. Expansion coefficients for the ground state energy.

| $U$ | $J/(t^2/U)$ | $e_n(U, J), n = 0 - 11$ |              |              |               |              |              |               |              |  |  |
|-----|-------------|-------------------------|--------------|--------------|---------------|--------------|--------------|---------------|--------------|--|--|
| 1   | 0.75        | -3.25                   | 3.0          | -0.64        | -0.5376       | -0.36882596  | -0.17599739  | -0.0037101999 | 0.11731865   |  |  |
|     |             | 0.17602728              | 0.17413953   | 0.11347540   | -0.0088559404 |              |              |               |              |  |  |
| 1   | 1.0         | -4.25                   | 4.0          | -0.5         | -0.4375       | -0.34273695  | -0.23157505  | -0.12102132   | -0.025358881 |  |  |
|     |             | 0.046126082             | 0.089325558  | 0.10390817   | 0.091603525   |              |              |               |              |  |  |
| 1   | 1.25        | -5.25                   | 5.0          | -0.41025641  | -0.36817883   | -0.30808094  | -0.23717612  | -0.16320812   |              |  |  |
|     |             | -0.093237945            | -0.032792472 | 0.014501216  | 0.046805520   | 0.063642701  |              |               |              |  |  |
| 4   | 0.75        | -1.75                   | 0.75         | -0.75294118  | -0.18602076   | 0.032575733  | -0.11521776  | 0.051494035   |              |  |  |
|     |             | 0.41270231              | 0.36649252   | -0.23825557  | -0.77368699   | -0.75051947  |              |               |              |  |  |
| 4   | 1.0         | -2.0                    | 1.0          | -0.69565217  | -0.21172023   | -0.030847663 | -0.060114049 | -0.011016385  |              |  |  |
|     |             | 0.16603971              | 0.22775615   | 0.035681298  | -0.22467766   | -0.32023724  |              |               |              |  |  |
| 4   | 1.25        | -2.25                   | 1.25         | -0.64646464  | -0.22854811   | 0.042172811  | -0.035358477 | -0.023047559  |              |  |  |
|     |             | 0.066640645             | 0.13361749   | 0.080005270  | -0.048764273  | -0.13678257  |              |               |              |  |  |
| 10  | 0.75        | -2.8                    | 0.3          | -0.38004751  | -0.018957239  | 0.066441099  | -0.087012637 | 0.038082237   |              |  |  |
|     |             | 0.16089588              | 0.11974079   | -0.075315493 | -0.24874869   | -0.20497804  |              |               |              |  |  |
| 10  | 1.0         | -2.9                    | 0.4          | -0.37383178  | -0.024456284  | -0.045510051 | -0.064243043 | 0.0053501507  |              |  |  |
|     |             | 0.075910665             | 0.074634170  | 0.0084898042 | -0.068321715  | -0.096846836 |              |               |              |  |  |
| 10  | 1.25        | -3.0                    | 0.5          | -0.36781609  | -0.029594398  | 0.033584424  | -0.050422135 | -0.0070731834 |              |  |  |
|     |             | 0.039086550             | 0.046846607  | 0.021090155  | -0.015763853  | -0.040974167 |              |               |              |  |  |
| 20  | 0.75        | -5.15                   | 0.15         | -0.19740901  | -0.0025574270 | 0.041371562  | -0.044384097 | 0.016313799   |              |  |  |
|     |             | 0.075735746             | 0.056531614  | -0.041034781 | -0.11826259   | -0.071871255 |              |               |              |  |  |
| 20  | 1.0         | -5.2                    | 0.2          | -0.19656020  | -0.0033806422 | -0.030305728 | -0.033253249 | 0.0010839733  |              |  |  |
|     |             | 0.034576709             | 0.03478823   | 0.019973437  | -0.033086550  | -0.037873386 |              |               |              |  |  |
| 20  | 1.25        | -5.25                   | 0.25         | -0.19571865  | -0.0041896960 | -0.023690877 | -0.02656723  | -0.0044976664 |              |  |  |
|     |             | 0.016991751             | 0.021350382  | 0.0088434956 | -0.0081635325 | -0.016372525 |              |               |              |  |  |

TABLE II. Expansion coefficients for the squared local moment.

| $U$ | $J/(t^2/U)$ | $l_n(U, J), n = 0 - 11$                                                                                                                           |
|-----|-------------|---------------------------------------------------------------------------------------------------------------------------------------------------|
| 1   | 0.75        | 1.0 0.0 -0.2048 -0.34406400 0.34363876 -0.18584371 0.073558744<br>0.3 2646019 0.45962201 0.39942169 0.126 77212 -0.33704683                       |
| 1   | 1.0         | 1.0 0.0 -0.125 -0.21875 -0.25342785 -0.21761601 -0.12058227<br>0.010355528 0.13808712 0.22622422 0.248 12902 0.19219824                           |
| 1   | 1.25        | 1.0 0.0 -0.084155161 -0.15104772 0.18802143 -0.18850833 -0.15340922<br>-0.090723264 -0.013648193 0.062253229 0.12199406 0.15344711                |
| 4   | 0.75        | 1.0 0.0 -0.28346021 -0.14006269 0.020099625 -0.042208513 0.13810651<br>0.6 6290690 0.47876441 -0.91546939 -1.99 197718 -1.29230064                |
| 4   | 1.0         | 1.0 0.0 -0.24196597 -0.147283644 0.0033966659 0.0020465901 0.041625973<br>0.27003095 0.33545751 -0.10812405 -0.68716227 -0.74045463               |
| 4   | 1.25        | 1.0 0.0 -0.20895827 -0.14774827 0.017014379 0.014267802 0.020573983<br>0.1 1999820 0.20351777 0.071247345 -0.22059729 -0.38236644                 |
| 10  | 0.75        | 1.0 0.0 -0.072218053 -0.0072046514 0.022443265 -0.034066355 0.041464469<br>0.1 1664774 0.058849733 -0.12141417 -0.25211532 -0.14201779            |
| 10  | 1.0         | 1.0 0.0 -0.069875098 -0.0091425362 -0.014415554 -0.024755587 0.016845440<br>0.060456748 0.045459350 -0.021365905 -0.086403185 -0.092929358        |
| 10  | 1.25        | 1.0 0.0 -0.067644339 -0.010885296 0.0099895925 -0.019054664 0.0064590300<br>0.035218951 0.032572341 0.0029017356 -0.031199252 -0.048202015        |
| 20  | 0.75        | 1.0 0.0 -0.019485158 -0.00050485912 0.0079729731 -0.0088647321 0.0094722236<br>0.027330357 0.013712645 -0.031211772 -0.058546614 -0.016716436     |
| 20  | 1.0         | 1.0 0.0 -0.019317955 -0.00066449970 -0.0057661935 -0.0066346164 0.0037510047<br>0.013829393 0.010568695 -0.0058676230 -0.020309374 -0.016500150   |
| 20  | 1.25        | 1.0 0.0 -0.019152896 -0.00082000166 -0.0044501566 -0.0052935510 0.013853661<br>0.078617712 0.0074609205 0.00033154323 -0.0074336626 -0.0092115763 |

TABLE III. Expansion coefficients for the sublattice magnetization.

| $U$ | $J/(t^2/U)$ | $m_n(U, J), n = 0 - 11$ |     |              |                |              |              |              |              |
|-----|-------------|-------------------------|-----|--------------|----------------|--------------|--------------|--------------|--------------|
| 1   | 0.75        | 1.0                     | 0.0 | -0.2048      | -0.34406400    | 0.37832834   | -0.30858617  | -0.16797349  | 0.0039694233 |
|     |             |                         |     |              |                | 0.17811940   | 0.32196832   | 0.35932350   | 0.14690771   |
| 1   | 1.0         | 1.0                     | 0.0 | -0.125       | -0.21875       | -0.26577581  | -0.26255537  | -0.21639940  | -0.14085987  |
|     |             |                         |     |              |                | -0.050839444 | 0.039513722  | 0.11520618   | 0.15738379   |
| 1   | 1.25        | 1.0                     | 0.0 | -0.084155161 | -0.15104772    | .19346230    | -0.20866262  | -0.19818728  | -0.16672332  |
|     |             |                         |     |              |                | -0.12065102  | -0.066877941 | -0.012424680 | 0.035149309  |
| 4   | 0.75        | 1.0                     | 0.0 | -0.28346021  | -0.14006269    | 0.29094791   | -0.70653657  | -0.11899194  | 1.78106763   |
|     |             |                         |     |              |                | 2.65518980   | 0.17150140   | -4.57860227  | -8.08441904  |
| 4   | 1.0         | 1.0                     | 0.0 | -0.24196597  | -0.14728364    | -0.16340074  | -0.36296753  | -0.23038035  |              |
|     |             |                         |     |              |                | 0.49230029   | 1.09523777   | 0.69655017   | -0.61481862  |
| 4   | 1.25        | 1.0                     | 0.0 | -0.20895827  | -0.14774827    | 0.11819956   | -0.21655277  | -0.20123752  | 0.10926874   |
|     |             |                         |     |              |                | 0.47800300   | 0.49583051   | 0.078098871  | -0.50046276  |
| 10  | 0.75        | 1.0                     | 0.0 | -0.072218053 | -0.0072046514  | 0.38636976   | -0.78484180  | -0.10195808  |              |
|     |             |                         |     |              |                | 1.63572360   | 2.29944530   | -0.28902064  | -4.69346784  |
| 10  | 1.0         | 1.0                     | 0.0 | -0.069875098 | -0.0091425362  | -0.21519753  | -0.44065350  | -0.20500969  |              |
|     |             |                         |     |              |                | 0.47694043   | 0.93375402   | 0.48462349   | -0.74441315  |
| 10  | 1.25        | 1.0                     | 0.0 | -0.067644339 | -0.010885296   | 0.13638406   | -0.28150839  | -0.18799630  |              |
|     |             |                         |     |              |                | 0.13527913   | 0.41216832   | 0.36759744   | -0.024358458 |
| 20  | 0.75        | 1.0                     | 0.0 | -0.019485158 | -0.00050485912 | 0.39342412   | -0.78866044  | -0.12416041  |              |
|     |             |                         |     |              |                | 1.59373845   | 2.17733968   | -0.58972155  | -4.77779001  |
| 20  | 1.0         | 1.0                     | 0.0 | -0.019317955 | -0.00066449970 | -0.22095641  | -0.44335646  | -0.21769894  |              |
|     |             |                         |     |              |                | 0.45233967   | 0.87748571   | 0.35719439   | -0.84087768  |
| 20  | 1.25        | 1.0                     | 0.0 | -0.019152860 | -0.00082000166 | -0.14118233  | -0.28358286  | -0.19603500  |              |
|     |             |                         |     |              |                | 0.11902603   | 0.38038377   | 0.30147500   | -0.092194212 |

TABLE IV. Expansion coefficients for the uniform magnetic susceptibility.

| $U$ | $J/(t^2/U)$ | $x_n(U, J), n = 0 - 9$ |      |             |              |             |             |              |              |
|-----|-------------|------------------------|------|-------------|--------------|-------------|-------------|--------------|--------------|
| 1   | 1.0         | 0.25                   | 0.25 | 0.17261905  | 0.027848648  | -0.1262274  | -0.19962669 | -0.11626869  |              |
|     |             |                        |      |             |              |             |             |              | 0.12849035   |
| 4   | 1.0         | 1.0                    | 1.0  | -0.67116442 | -2.82447457  | -1.58009047 | 4.43513119  | 8.64219154   |              |
|     |             |                        |      |             |              |             |             |              | 1.42302418   |
| 6   | 1.0         | 1.5                    | 1.5  | -1.48149027 | -4.91068070  | -1.92590553 | 8.56642180  | 13.73914475  |              |
|     |             |                        |      |             |              |             |             |              | -1.78852254  |
| 10  | 1.0         | 2.5                    | 2.5  | -2.99306647 | -8.80889678  | -2.67462512 | 16.12944762 | 23.047960372 |              |
|     |             |                        |      |             |              |             |             |              | -8.54214685  |
| 20  | 1.0         | 5.0                    | 5.0  | -6.48980810 | -18.15418203 | -4.79124584 | 33.98201522 | 45.62214165  |              |
|     |             |                        |      |             |              |             |             |              | -23.66505149 |

TABLE V. Expansion coefficients for the “paramagnetic” part ( $\rho_s^{para}$ ) of the spin stiffness.

| $U$ | $J/(t^2/U)$ | $p_n(U, J), n = 0 - 9$                                                                             |
|-----|-------------|----------------------------------------------------------------------------------------------------|
| 1   | 1.0         | 0.0 0.0 0.0 0.0 0.000047115128 0.00023699734 0.000655001<br>0.0012665331 0.0018294204 0.0019236148 |
| 4   | 1.0         | 0.0 0.0 0.0 0.0 0.0089710818 0.028888083 0.029891453<br>-0.020672065 -0.090109640 -0.087959801     |
| 10  | 1.0         | 0.0 0.0 0.0 0.0 0.012111686 0.027777667 0.018651461<br>-0.018905322 -0.049016789 -0.033652011      |
| 20  | 1.0         | 0.0 0.0 0.0 0.0 0.0076518484 0.015940557 0.010548308<br>-0.0087795777 -0.023213629 -0.013514588    |

## Figure Captions

Fig. 1 The ground state energy as a function of  $U$ . Filled circles are our calculated data, with a solid line as guide to the eye. Errorbars indicate spread of the Padé approximants. Here  $E_g + U/4 = \text{Kinetic Energy} + U \langle n_\uparrow n_\downarrow \rangle$ .

Fig. 2 The squared local moment  $L$  as a function of  $U$ . Filled circles are our calculated data, with a solid line as guide to the eye. Errorbars indicate spread of the Padé approximants. Open squares are the results of Monte Carlo simulation by White *et al.* [10], with dotted line as guide to the eye. The dashed line is the result of the mean field solution.

Fig. 3 Sublattice magnetization  $2M^+$  as a function of  $U$ . Filled circles are our calculated data, with a solid line as guide to the eye. Errorbars indicate spread of the Padé approximants. Long dashed line and dotted line represent mean field solution and RPA result respectively. A short dashed line indicates the limiting large  $U$  (Heisenberg) value.

Fig. 4 (a) The ratio of the uniform magnetic susceptibility to  $U$  as a function of  $U$ . Filled circles are our calculated data, with a dashed line as guide to the eye. A short dashed line indicates the limiting large  $U$  (Heisenberg) value. (b) The uniform magnetic susceptibility as a function of  $U$ . Filled circles are our calculated data, with a dashed line as guide to the eye. A crossover between SDW antiferromagnetic state and Heisenberg antiferromagnet is around  $U_c \approx 4$ .

Fig. 5 (a) The product of the spin stiffness and  $U$  as a function of  $U$ . Filled circles are our calculated data, with a solid line as guide to the eye. A short solid line indicates the limiting large  $U$  (Heisenberg) value. (b) The spin stiffness as a function of  $U$ . Filled circles are our calculated data, with a solid line as guide to the eye. Errorbars indicate spread of the Padé approximants. The solid line is the mean field result. The filled squares are results of variational Monte Carlo simulations [21].

Fig. 6 The spin wave velocity as a function of  $U$ . Filled circles are our calculated data, with a solid line as a guide to the eye. The dotted line and dashed line represent results of RPA and Heisenberg model respectively.